

gtt_ekn Manual

Edition : 2016.03.28

Y.Goto, Y.Tachibana, N.Takayama

1 Function Manual

1.1 Outline

1.2 Notation

1.3 Functions

1.3.1 gtt_ekn.nc

`gtt_ekn.nc(beta,p)`

:: It returns the value of the normalizing constant and its derivatives of the conditional hypergeometric distribution of two way contingency tables with fixed marginals

return $[Z, [d_{11} Z, d_{12} Z, \dots], \dots, [d_{m1} Z, d_{m2} Z, \dots, d_{mn} Z]]$

beta List of row sums and column sums. All entries must be positive integer.

p The probabilities of the cells of the table.

- Details have not been written. See Japanese documents.

2x3 contingency table with the row sum [4,5] and the column sum [2,4,3].

```
[2237] gtt_ekn.nc([[4,5],[2,4,3]],[[1,1/2,1/3],[1,1,1]]);
[4483/124416,[ 353/7776 1961/15552 185/1728 ]
[ 553/20736 1261/15552 1001/13824 ]]
```

English translation for other functions has not yet been written.

Index

(Index is nonexistent)

(Index is nonexistent)

Short Contents

1	Function Manual	1
	Index	2

Table of Contents

1	Function Manual	1
1.1	Outline	1
1.2	Notation	1
1.3	Functions	1
1.3.1	gtt_ekn.nc	1
Index		2